

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.


**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

EP 99
2930
D

PCT/EP 99/02930
11 May 1999

REC'D 03 JUN 1999
WIPO PCT

5



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

April 26, 1999

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/083,450

FILING DATE: April 29, 1998

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

P. R. Grant

P. R. GRANT
Certifying Officer

Please type a plus sign (+) inside this box → ☐

Docket Number 1038-820 M13-jb

PROVISIONAL APPLICATION FOR PATENT COVER SHEET (Large Entity)

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c).

71479 U.S. PTO



00652410 0545009

INVENTOR(S)/APPLICANT(S)					
Given Name (first and middle [if any])		Family Name or Surname		Residence (City and either State or Foreign Country)	
Andreas		Böble		Fasanenring 2, D-23627 Gross Grönau, Germany.	
Dieter		Jocham		Zwinglistrasse 1, D-23568 Lübeck, Germany.	
<input type="checkbox"/> Additional inventors are being named on page 2 attached hereto					
TITLE OF THE INVENTION (280 characters max)					
TREATMENT OF PAPILLOMA VIRUS INFECTION					
CORRESPONDENCE ADDRESS					
Direct all correspondence to:					
<input type="checkbox"/> Customer Number				Place Customer Number Bar Code Label here	
OR					
<input checked="" type="checkbox"/> Firm or Individual Name		Michael I. Stewart			
Address		Sim & McBurney			
Address		6th Floor, 330 University Avenue			
City		Toronto		State	Ontario
Country		Canada		ZIP	M5G 1R7
		Telephone	(416) 595-1155	Fax	(416) 595-1163
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification		Number of Pages		13	
<input type="checkbox"/> Drawing(s)		Number of Sheets		<input type="checkbox"/> Other (specify)	
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (check one)					
<input checked="" type="checkbox"/> A check or money order is enclosed to cover the filing fees				FILING FEE AMOUNT	
<input type="checkbox"/> The Commissioner is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number.				\$150.00	
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government					
<input checked="" type="checkbox"/> No					
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are.					

Respectfully submitted,

SIGNATURE Michael I. Stewart

DATE April 28, 1998

TYPED or PRINTED NAME Michael I. Stewart

REGISTRATION NO. 24,973
(if appropriate)

TELEPHONE (416) 595-1155

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

SEND TO: Box Provisional Application, Assistant Commissioner for Patents, Washington, DC 20231



TITLE OF INVENTION

TREATMENT OF PAPILLOMA VIRUS INFECTION

FIELD OF INVENTION

5 The present invention relates to the treatment of disease caused by papilloma virus infection, in particular to the adjuvant treatment of condylomata acuminata caused by human papilloma virus in men.

BACKGROUND OF THE INVENTION

10 Human papilloma virus (HPV) infections of the urogenital tract represent the most often sexually transmitted viral disease in man (refs. 1 to 3 - various references are referred to in parenthesis to more fully describe the state of the art to which this
15 invention pertains. Full bibliographic information for each citation is found at the end of the specification, immediately preceding the claims. The disclosure of these references are hereby incorporated by reference into the present disclosure). HPV is a double stranded
20 DNA virus and with the recent developed molecular biological techniques, more than 55 different HPV types have been recognized (ref. 4). HPV is associated with a wide spectrum of clinical states including condylomata acuminata, latent and subclinical infection, and
25 Bowen's disease. Subclinical infections gain more importance as they are believed to cause intraepithelial neoplasia, based on the frequent detection of HPV DNA in invasive carcinomas, especially in urogenital region (refs. 1, 5). A significant risk
30 for the development of an invasive cancer is ascribed to the infections by HPV 16, 18 and 33 (refs. 6 to 9).

 The most prevalent HPV types causing condylomata acuminata are type 6 and 11. Condylomata acuminata are visible, multifocal, multicentric and multiform

50083450-012998

lesions. Predilection sites are penis, scrotum, perineum, urethra, perianal regions, intertriginous zones, and oral mucosa. In uncircumcised men the frenulum, the coronary sulcus and the inner aspect of the foreskin are most often afflicted, whereas in circumcised patients the shaft of the penis is involved. Genital warts are of great psychological and cosmetic relevance representing a major hindrance to sexual performance.


Treatment options include surgical methods like excision, electrocautery, cryosurgery or laser vaporization. It has been shown in molecular hybridization studies that HPV DNA sequences do exist in adjacent normal tissue after carbon dioxide laser removal of genital warts (ref. 10). These findings and the well known high recurrence rates after initial treatment demonstrate the need for adjuvant therapy to eradicate invisible disease. Therapeutic results with local application of cytotoxic agents, for example, 5-fluorouracil and podophyllin/podophyllotoxin have, however, been unsatisfactory (refs. 11 to 13). Furthermore, several types of interferons (IFN) as well as autologous vaccines have been tried with varying success (refs. 2, 14 to 17). More recently, oral isotretinoin has been given with some success to reduce the recurrence rate (refs. 18, 19).

SUMMARY OF INVENTION

In accordance with one aspect of the present invention, there is provided a method of treatment of disease caused by papilloma virus, which comprises applying an effective amount of *Mycobacterium* to the region of infection.

In particular, the present invention in accordance with the treatment of infections caused by human papilloma virus (HPV) using the *Mycobacterium*. Such

13063450.04298



infection may include cutaneous and genital warts in man, including verruca vulgaris and condyloma acuminatum, cervical intraepithelial neoplasia and genital carcinomas. The present invention is
5 illustrated by application to recurrent external and intraurethral condylomata acuminata in men.

The *Mycobacterium* which may be employed may be an attenuated form of a *Mycobacterium* of the tuberculosis complex and, in particular, may be an
10 attenuated form of *M. bovis*, specifically Bacillus Calmette-Guerin (BCG).

The treatment may be effected by application of the *Mycobacterium* in a suitable carrier to the region of infection, which may involve topical application to
15 penile and perianal areas, or intraurethral application to the urogenital tract. The treatment may involve a single or a plurality of doses applied at time intervals. The individual dosage level may be about 1 mg to about 500 mg while the time interval between
20 doses may vary from about 1 to about 30 days. The number of treatments applied is from 1 to about 30 treatments. The treatment may be preceded by laser or other surgical or topical therapy.

The present invention extends, in a further aspect
25 thereof, to the use of a *Mycobacterium* in the manufacture of a medicament for the treatment of infection caused by papilloma virus, particularly human papilloma virus.

GENERAL DESCRIPTION OF INVENTION

30 A wide spectrum of therapeutic approaches has been used over the past years for the management of condylomata acuminata. However, no form of therapy has yielded consistently effective results. In order to avoid recurrence of disease, the combination of

60083450 042999

surgical ablative methods and immunomodulative agents seem to be promising. Due to their antiviral, immunomodulative and antiproliferative properties, interferons have been given as a topical and systemic
5 form of treatment for condylomata acuminata. Interferons were also chosen for their mild side effect profile but recurrence rates up to 75% have been reported (refs. 2, 14 to 17).

In the case studies reported below, patients
10 treated with BCG were cleared of warts and cytological evidence of HPV and had no recurrence of disease. These patients have previously experienced between two and five recurrences of condylomata acuminata. The annual recurrence rate was significantly reduced from
15 3.2 with standard therapy to 0.75 with BCG therapy. The incidence of side effects due to BCG therapy was low. One patient reported mild dysuria after the second and third BCG application of this first treatment course. Another patient had a penile edema and fever which was managed conservatively. Penile
20 edema is a rare complication after intravesical BCG instillation. However, with intraurethral instillation of BCG for transitional cell carcinoma of the urethra has been reported by Baniel et al (ref. 20). Serious
25 complications like sepsis or hematuria were not observed in the reported group.

EXAMPLE

The above disclosure generally describes the present invention. A more complete understanding can be
30 obtained by reference to the following specific Example. This Example are described solely for purposes of illustration and are not intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may
35 suggest or render expedient. Although specific terms

00083450-042009

have been employed herein, such terms are intended in a descriptive sense and not for purposes of limitation.

Between October 1994 and March 1997, six men with rapidly recurrent external and intraurethral condylomata acuminata were selected for BCG therapy. The mean age was 27 years (range 22 to 32 years). All patients had previously undergone therapy for recurrent condylomata acuminata in other dermatological and urologic departments (see Table 1 below). The patients were informed they were undergoing an experimental treatment approach and all individuals gave written informed consent.

Examination of the penis and scrotum was performed without and with acetoacid application and lesions were treated with the neodymium:yttrium-aluminum-garnet (Nd:YAG) laser (Dornier Medilas Fibertome 4060) at 10 to 20 W. Endoscopic inspection of the urethra and bladder was also performed and visible lesions treated with the Nd:YAG laser. Finally, the perianal region was inspected and laser treatment, when necessary, applied.

A minimum interval of 3 weeks between laser therapy and BCG application was utilized. Internal application for BCG preparation, 81 mg of Connaught strain (Pasteur Mérieux Connaught Canada, North York, Ontario, Canada) were dissolved in 2 ml. Of sterile saline. The solution was directly instilled into the urethra with low pressure and kept for a minimum of 2 hours. To avoid preliminary emptying, a dressing was taped over the meatus. After 2 hours, the dressing was removed and the patient allowed to void spontaneously. A total of six BCG instillations were given in weekly intervals.

External application, eighty-one mg of BCG Connaught strain were dissolved in 2 ml of sterile saline. The solution was given onto a dressing which was gently wrapped around the sulcus and other affected

areas. The dressing was fixed and kept for 2 hours as a moist chamber. This procedure was also repeated six times in weekly intervals.

The courses of six patients are given below and in

5 Table 1 below:

Case 1

A 22-year-old male patient developed condylomata acuminata in 1994. Initially, the frenulum and the meatus were involved. He was treated with local
10 podophyllin but relapsed twice, and was referred to the inventors department in April 1995. Over a period of 13 months, five relapses occurred despite internal and external laser application. From July until October 1996, six intraurethral instillations and external
15 applications of BCG were given in weekly intervals. At follow-up of 11 months, no relapse had been noticed.

Case 2

A 28-year-old male patient demonstrated with perianal and preputial condylomata in May 1995.
20 Cystoscopy revealed additional lesions of the urethra and, therefore, internal and external laser therapy was performed. Three months later an urethral relapse was diagnosed and again treated with laser. Subsequently, the patient received six intraurethral instillations
25 combined with external application of BCG. At follow-up of 29 months, no relapse had been noticed.

Case 3

A 26-year-old male patient had initial diagnosis of condylomata acuminata in 1992. The affected sites
30 included the meatus and frenulum as well as the perianal region. Over a period of 15 months, four relapses occurred which were all managed with laser therapy. In September 1995, intraurethral instillation and external application of BCG was performed. At follow-up of 24
35 months, no relapse had been noticed.

Case 4

A 32-year-old man developed condylomata acuminata in February 1995. He had a hypospadias without operative correction required. Initial involvement included the meatus, subsequently frenulum and urethra had been involved. During the next 16 months, four relapses were diagnosed and treated with laser. Intraurethral BCG was given in August and September 1996. After 2 further relapses, intraurethral instillation and external application of BCG was repeated in January and February 1997. At follow-up of 8 months, no relapse had been noticed.

Case 5

A 31-year-old male had initial diagnosis of condylomata acuminata in January 1993. Primarily, meatus and foreskin were affected. In the next 27 months, four relapses occurred and managed with laser therapy. The first course of intraurethral instillation and external BCG application was performed in May and June 1995. The patient had three relapses which were treated with laser. In June and July 1996, he received a second course of BCG. He demonstrated one relapse which was lasercoagulated and at follow-up of 12 months, no further relapse had occurred.

Case 6

A 27-year-old male patient demonstrated condylomata acuminata in August 1996. Penis, meatus and urethra were involved. After two relapses, BCG was applied in July 1997. Due to side effects, the treatment was stopped after the third instillation. He had demonstrated a penile edema and fever (38.5°C) and was treated with antiphlogistic drugs and ice packages. At follow-up of 3 months, one new penile lesion had developed.

In summary of the clinical results obtained, three patients completed one course of BCG. These patients

856240-042998

had no relapse in follow-up studies (Table 1). Two patients underwent a second course of BCG after recurrence of condylomata acuminata. One of them remained without relapse after the second course in a follow-up period of 8 months. The other patient had one relapse after the second BCG course. One patient developed a relapse after he had discontinued therapy due to side effects following the third BCG instillation of his first course.

With respect to side effects, one patient reported mild dysuria after the second and third BCG instillation of his first treatment course. Another patient was withdrawn from BCG therapy due to penile edema and fever. No other side effects were reported.

Before BCG therapy, 36 recurrences of condylomata acuminata were noted in 136 observation months compared to 6 episodes during 96 months after treatment. Therefore, the annual recurrence rate before BCG therapy was 3.2 and 0.75 thereafter. These results were statistically significant ($p < 0.05$, test of equality of 2 percentages, (ref. 21).

SUMMARY OF DISCLOSURE

In summary of this disclosure, BCG therapy is useful in treating condylomata acuminata, particularly rapidly recurrent conditions. In general, *Mycobacteria* are useful in treating infections caused by papilloma virus. Modifications are possible within the scope of this invention.

0003450 04293

356240 054809

TABLE 1

AGE	No. previous occurrences	No. relapses before BCG	No. relapses after BCG	Localisation of relapses before BCG	Localisation of relapses after BCG
22	3	5	0	Urethra, frenulum, meatus foreskin, perianal	
28	1	2	0		
26	5	4	0	frenulum, urethra, meatus	
32	1	4	2,0 after 2 nd course	frenulum, urethra, urethra	frenulum, urethra
27	3	4	3,0 after 2 nd course	foreskin, meatus urethra	foreskin, meatus urethra
27	1	3	1 (incomplete course)	meatus, urethra	meatus, urethra

[illegible]

1. Syrjänen K.J.: HPV in genital squamous cell tumors: epidemiology and clinical synopsis. In: Genital Papillomavirus Infections. Edited by G. Gross, S. Jablonska, H. Pfister and H. E. Stegner. Berlin Heidelberg Now York: Springer-Verlag, pp. 3-12, 1990.
2. Bonnez W., Oakes, D., Bailey-Farchione, A., Choi, A., Hallahan, D., Pappas, P., Holloway, M., Corey, L., Barnum, G., Dunne, A., Stoler, M.H., Demeter, L.M. and Reichman, R.C.: A randomized, double-blind, placebo-controlled trial of systemically administered interferon- α , β , or γ in combination with cryotherapy for the treatment of condyloma acuminatum. J. Infect. Dis., 171:1081, 1995
3. Grin, W.: Condyloma acuminata-Epidemiologie und Therapie. Wien. Klin. Wochenschr., 104:215, 1992.
4. Barrasso, R., de Brux, J., Croissant, O. and Orth, G.: High prevalence of papillomavirus-associated penile intraepithelial neoplasia in sexual partners of women with cervical intraepithelial neoplasia. N. Engl. J. Med., 317: 916, 1987.
5. von Knebel Doeberitz, M. and zur Hausen, H.: Biological significance of human papillomavirus early gene expression in human cervical carcinoma cells. In: Genital Papillomavirus Infections. Edited by G. Gross, S. Jablonska, H. Pfister and H.E. Stegner. Berlin Heidelberg Now York: Springer-Verlag, pp.51-65, 1990.
6. Smith, K.T. and Campo, M.S.: Papillomaviruses and their involvement in oncogenesis. Biomed. Pharmacoth., 39: 405, 1985.
7. Pfister, H.: Biology and biochemistry of papillomaviruses. Rev. Physiol. Biochem. Pharmacol., 99: 111, 1984.
8. Syrjänen, S.M., von Krogh, G. and Syrjänen, K.J.: Detection of human papillomavirus DNA in anogenital condylomata in men using in situ DNA hybridisation applied to paraffin sections. Genitourin. Med., 63: 31. 1987.
9. Wiener, J.S., Effert, P.J., Humpphrey, P.A., Yu, L., Liu, E.T. and Walter, P.J.: Prevalence of human papilloma virus types 16 and 18 in squamous-cell carcinoma of the penis: a retrospective analysis of primary and metastatic sections by differential polymerase chain reaction. Int. J. Cancer, 50:694, 1992.

10. Ferenczy, A., Mitao, M., Nagai, N., Silverstein, S.J. and Crum, C.P.: Latent papillomavirus and recurrent genital warts. *N. Engl. J. Med.*, 313: 784, 1986.
11. Swinehart, J.M., Sperling, M., Phillips, S., Kraus, S., Gordon, S., McCarty J.M., Webster G.F., Skinner, R., Korey, A. and Orenberg, E.K: Intralesional fluorouracil/epinephrine injectable gel for treatment of condylomata acuminata. A phase 3 clinical study. *Arch. Dermatol.*, 133: 67, 1997.
12. von Krogh, G., Szpak E., Andersson. M. and Bergelin, I.: Self-treatment using 0.25%-0.50% podophyllotoxin-ethanol solutions against penile condylomata acuminata: a placebo-controlled comparative study. *Genitourin. Med.*, 70: 105, 1994.
13. Syed, T.A. and Lundin, S.: Topical treatment of penile condylomata acuminata with podophyllotoxin 0.3% solution, 0.3% cream and 0.15% cream. A comparative open study. *Dermatology*, 187: 30, 1993.
14. Davis, B.E. and Noble, M.J.: Initial experience with combined interferon-alpha 2 B and carbon dioxide laser for the treatment of condylomata acuminata. *J. Urol.*, 147: 627, 1992.
15. Gross, G., Roussaki, A., Baur, S., Wiegand, M. and Mescheder, A.; Systemically administered interferon alfa-2a prevents recurrence of condylomata acuminata following CO₂ laser ablation. The influence of the cyclic low-dose therapy regime. Results of a multicentre double-blind placebo-controlled clinical trial. *Genitourin. Med.*, 72:71, 1996
16. Monsonogo, J., Cessot, G., Ince, S.E., Galazka. A.R and Abdul-Ahad, A.K.: Randomised double-blind trial of recombinant interferon-beta for condyloma acuminatum. *Genitourin. Med.*, 72: 111, 1996.
17. Cardamakis, E., Kotoulas, I.-G., Metalinos, K, Mantouvalos, H., Relakis, K, Scapari, M. Korantzis, and Papathanasiou, Z.: Treatment of recurrent condylomata acuminata or flac condylomata with interferon-α2a. *J. Vol.* 152:2011, 1994.
18. Cardamakis, E.K. Kotoulas, I-G.B., Dimopoulos, D. P., Stathopoulos, E.N. Michopoulos, J.T. and Tzingounis, V.A: Comparative study of systemic interferon alfa-2a with oral isotretinoin and oral isotretinoin alone in the treatment of recurrent condylomata acuminata. *Arch. Gynecol. Obstet.*, 258: 35, 1996.
19. Tsambaos, D., Geogiou, S., Monastini, A., Sakkis, T., Sagriotis, A. and Georz G.: Treatment of condylomata

11.05.99

12

acuminattta with oral isotretionin. J. Urol, 158: 1810, 1997

20 Baniel, J., Lev, Z., Engelstein, D. and Servadio, C.: Penile edema and meatal ulceration after intravesical instillation with bacillus Calmette-Guerin. Urology, 47: 932, 1996.

21. Sokal, R.R. and Rohlf, F.J.: Test of equality of 2 percentages In: Biometry W.H. Freeman Company, San Francisco, p.p. 607 to 608, 1969.

0003450-042990

ABSTRACT OF THE DISCLOSURE

Disease caused by papilloma virus is treated by applying an effective amount of *Mycobacterium* to the region of infection. Specifically, condylomata acuminata are caused by human papilloma virus infection. Despite numerous treatment modalities these patients often demonstrate recurrent disease. BCG therapy is used in patients not responding to standard treatment. Six men with rapidly recurrent external and intraurethral condylomata acuminata underwent BCG therapy after initial laser treatment. External application and intraurethral instillation of BCG was performed six times in weekly intervals. Follow-up studies included examination and endoscopic inspection of the urethra and bladder. Three patients completed one course of BCG and had no relapse of condylomata acuminata. Two patients underwent a second course of BCG, of whom one relapsed. One patient relapsed after discontinued therapy due to penile edema. The annual recurrence rate decreased from 3.2 before BCG therapy to 0.75 after BCG therapy ($p < 0.05$, test of equality of 2 percentages). The BCG-induced immune response appears to reduce the recurrence rate in patients with condylomata acuminata.